

A reconstruction problem related to balance equations-I *

Bhalchandra D. Thatte

bdthatte@gmail.com

Mathematics Subject Classifications: 05C60

Abstract

A modified k -deck of a graph is obtained by removing k edges in all possible ways and adding k (not necessarily new) edges in all possible ways. Krasikov and Roditty used these decks to give an independent proof of Müller's result on the edge reconstructibility of graphs. They asked if a k -edge deck could be constructed from its modified k -deck. In this paper, we solve the problem when $k = 1$. We also offer new proofs of Lovász's result, one describing the constructed graph explicitly, (thus answering a question of Bondy), and another based on the eigenvalues of Johnson graph.

1 Introduction

Graphs in this paper are assumed to be undirected and without multiedges or loops, and are assumed to have n vertices and m edges, unless specified otherwise. The complement of G is denoted by G^c . We set $N = \binom{n}{2}$.

For a graph G , we define three kinds of decks – $MD_i(G)$, $PD_i(G)$ and $ED_i(G)$ as follows. The deck $MD_i(G)$, called the modified i -deck in [KR], is obtained by removing i edges of G in all possible ways and adding i (not necessarily new) edges in all possible ways. When we insist that the set of added edges and the set of removed edges be disjoint, we call it the perturbed i -deck, and denote it by $PD_i(G)$. The deck $ED_i(G)$ is the i -edge deck, i.e., the collection of all the subgraphs obtained by deleting i edges. We similarly define the above decks for a collection $S = \{G_1, G_2, \dots, G_r\}$

*Published in Discrete Mathematics 176 (1997) 279-284.

of graphs as the multiunion of the corresponding decks for all the members of the collection. Thus, e.g., $MD_i(S) = \cup_j MD_i(G_j)$, where $\cup$ denotes a multiunion.

In this paper, we denote sets (or multisets) by their characteristic vectors of length equal to the number of nonisomorphic graphs on n vertices and m edges, (or $m - i$ edges in case of the i -edge deck). The characteristic vector of a set P of graphs is denoted by X_P , and that of the singleton set $\{G\}$ by simply X_G . The operations of constructing the above decks are represented by matrices whose rows are indexed by all the nonisomorphic graphs on n vertices and m edges, ($m - i$ edges when we are considering i -edge decks), and whose columns are indexed by nonisomorphic graphs on n vertices and m edges. We denote the matrices corresponding to MD_i , PD_i and ED_i respectively by Δ_i , D_i and d_i . The notation chosen here for denoting the matrices is same as the notation used in [KR] for the corresponding decks. The kl -th entry of Δ_i is the number of graphs isomorphic to G_k in $MD_i(G_l)$. The kl -th entry of D_i is the number of graphs isomorphic to G_k in $PD_i(G_l)$. The kl -th entry of d_i is the number of graphs isomorphic to G_k in $ED_i(G_l)$. Note that matrices D_0 and Δ_0 are identity operators. We derive various useful identities involving linear combinations of these matrices or polynomials in the matrices.

Krasikov and Roditty [KR] used the modified decks to set up their balance equations, which they used to derive Müller's result independently. It is easy to see that $MD_i(G)$ is constructed from $ED_i(G)$ by simply adding i edges in all possible ways in each graph in $ED_i(G)$. They asked if $ED_i(G)$ could be constructed from $MD_i(G)$. This problem is solved for the case when $i = 1$. Also, a new proof of Lovász's result based on the eigenvalues of Johnson graph is obtained.

2 Reconstructing ED_1 from MD_1 .

We first derive some identities which various matrix operators satisfy.

Lemma 1. $\Delta_k = \sum_{i=0}^k \binom{m-i}{k-i} D_i$

Proof This equivalent to Lemma 3.1 in [KR].

Theorem 2.

$$D_1 D_i = (m - i + 1)(N - m - i + 1) D_{i-1} + i(N - 2i) D_i + (i + 1)^2 D_{i+1}$$

Proof For a graph G , consider a typical member $G - X + Y$ of $PD_i(G)$, where $X \cap Y = \phi$. For an edge $e \in E(G) - X + Y$ and an edge $f \notin E(G) - X + Y$, there is a graph $H = G - X + Y - e + f$ in $PD_1(PD_i(G))$. Depending upon the choice of e and f , following four cases arise.

1. $e \in Y$ and $f \in X$: $H \in PD_{i-1}(G)$.
2. $e \in E(G) - X$ and $f \in X$: $H \in PD_i(G)$.
3. $e \in Y$ and $f \in E(G^c) - Y$: $H \in PD_i(G)$.
4. $e \in E(G) - X$ and $f \in E(G^c) - Y$: $H \in PD_{i+1}G$.

In case 1, same H would be obtained if we first replaced $X' = X - f + f'$ by $Y' = Y - e + e'$, where $f \neq f'$ and $e \neq e'$, and then e' by f' . There are $(m - i + 1)$ choices for f' and $(N - m - i + 1)$ for e' . This explains the first term on the right hand side. In a similar manner, the second term results from cases 2 and 3, and the third term from case 4.

Theorem 2 is the basis of our procedure of relating Δ_i to Δ_1 . We describe the procedure by first relating Δ_2 to Δ_1 . We apply Theorem 2 for $i = 1$. That allows us to express D_2 in terms of D_0 , D_1 , and D_1D_1 . Since, D_1 can be expressed in terms of Δ_1 and D_0 , we can express Δ_2 in terms of D_0 , Δ_1 and Δ_1^2 . It is then enough to demonstrate that the expression for Δ_2 obtained this way has no D_0 term. From Lemmas 1 and Theorem 2, we write

$$\begin{aligned}
\Delta_2 &= \binom{m}{2} + \binom{m-1}{1} D_1 + \binom{m-2}{0} D_2 \\
&= \frac{m(m-1)}{2} D_0 + (m-1)(\Delta_1 - mD_0) \\
&\quad + \frac{1}{4} \{(\Delta_1 - mD_0)(\Delta_1 - mD_0) - m(N-m)D_0 - (N-2)(\Delta_1 - mD_0)\} \\
&= \frac{2(m-1) - N}{4} \Delta_1 + \frac{1}{4} \Delta_1^2
\end{aligned}$$

To relate Δ_i , $i > 1$ to Δ_1 , we first prove a lemma which relates D_i to Δ_1 and D_0 , and then use Lemma 1.

Lemma 3. D_k can be written as

$D_k = L_k + (-1)^k \binom{m}{k} D_0$ where L_k is a linear combination of operators constructed from Δ_1 .

Proof We prove this by induction on k . For $k = 0$ and $k = 1$, the result is trivial to prove. Let $D_k = L_k + (-1)^k \binom{m}{k} D_0$ for $k \leq r$. To prove the result for $k = r + 1$, we operate from left by D_1 on both sides of $D_r = L_r + (-1)^r \binom{m}{r} D_0$. From Theorem 2

and the induction hypothesis we get,

$$\begin{aligned}
& D_1 \left\{ L_r + (-1)^r \binom{m}{r} D_0 \right\} \\
= & (m - r + 1)(N - m - r + 1) \left\{ L_{r-1} + (-1)^{r-1} \binom{m}{r-1} D_0 \right\} \\
& + r(N - 2r) \left\{ L_r + (-1)^r \binom{m}{r} D_0 \right\} + (r + 1)^2 D_{r+1}
\end{aligned}$$

Since $D_1(D_0) = D_1 = \Delta_1 - mD_0$, we have

$$\begin{aligned}
& (r + 1)^2 D_{r+1} \\
= & D_1(L_r) + (-1)^r \binom{m}{r} \Delta_1 - (m - r + 1)(N - m - r + 1)L_{r-1} \\
& - r(N - 2r)L_r - m(-1)^r \binom{m}{r} D_0 \\
& - (m - r + 1)(N - m - r + 1)(-1)^{r-1} \binom{m}{r-1} D_0 \\
& - r(N - 2r)(-1)^r \binom{m}{r} D_0
\end{aligned}$$

From the induction hypothesis, we claim that first four terms on the right side give a linear combination of operators constructed from Δ_1 . Also, coefficient of D_0 on the right side simplifies to $(-1)^{r+1}(r + 1)^2 \binom{m}{r+1}$. Thus $D_{r+1} = L_{r+1} + (-1)^{r+1} \binom{m}{r+1} D_0$, which completes the proof.

Theorem 4. For $i \geq 1$, Δ_i can be written in terms of Δ_1

Proof In the expression for Δ_i given by Lemma 1, we substitute D_k from Lemma 3, Then it is enough to prove that the the coefficient of D_0 in the resulting expression

is zero.

$$\begin{aligned}
\Delta_i &= \sum_{k=0}^i \binom{m-k}{i-k} D_k \\
&= \sum_{k=0}^i \binom{m-k}{i-k} \{(-1)^k \binom{m}{k} D_0 + L_k\} \\
&= \sum_{k=0}^i \binom{m-k}{i-k} (-1)^k \binom{m}{k} D_0 + \sum_{k=0}^i L_k \\
&= \sum_{k=0}^i L_k
\end{aligned}$$

Lemma 5. *Let $m \geq N/2$, and let P and Q be collections of graphs such that $d_1 X_P = d_1 X_Q$ or $\Delta_1 X_P = \Delta_1 X_Q$, then $X_P = X_Q$.*

Proof Since modified decks can be obtained from edge decks, we assume $\Delta_1 X_P = \Delta_1 X_Q$. From Theorem 4, $\Delta_k X_P = \Delta_k X_Q$ for $k \geq 1$. Now the claim in the lemma is an immediate consequence of the fact that the version of Lovász's result presented in Remark 1 in the next section makes use of the modified decks only. We also note that Krasikov and Roditty used only the modified decks in their proof of Müller's result.

Now we prove the main result of this section.

Theorem 6. *For collections P and Q of graphs, if $\Delta_1 X_P = \Delta_1 X_Q$ then $d_1 X_P = d_1 X_Q$.*

Proof Let $P' = \{F^c; F \in ED_1(P)\}$ and $Q' = \{F^c; F \in ED_1(Q)\}$. Note that $\Delta_1 X_P = \Delta_1 X_Q$ is equivalent to $d_1 X_{P'} = d_1 X_{Q'}$, (where d_1 corresponds to the operation of constructing edge decks of n -vertex and $(N - m + 1)$ -edge graphs): this follows from the fact that for any F , $e \in E(F)$ and $f \notin E(F) - e$, $(F - e + f)^c = (F - e)^c - f$. Now, if $m > N/2$ then $X_P = X_Q$, therefore, $d_1 X_P = d_1 X_Q$. Else, $N - m + 1 > N/2$, implying $X_{P'} = X_{Q'}$, therefore, $d_1 X_P = d_1 X_Q$.

Corollary 7. *Given the modified 1-deck of a graph, its 1-edge deck can be constructed.*

Remarks

1. In his 1977 survey, Bondy asked if Lovász's result could be proved constructively, (see problem 12 in [B]). The inversion of the equation in Lemma 1 gives

$$D_k = \sum_{i=0}^k (-1)^{k+i} \binom{m-i}{k-i} \Delta_i$$

Therefore, when $m > N/2$, D_m is zero. Therefore,

$$\Delta_0 = \Delta_1 - \Delta_2 + \Delta_3 - \dots + (-1)^{m+1} \Delta_m$$

and,

$$X_P = \Delta_0 X_P = \{\Delta_1 - \Delta_2 + \Delta_3 - \dots + (-1)^{m+1} \Delta_m\} X_P$$

for any collection P of graphs.

Thus we explicitly know the reconstructed graph or a collection of graphs.

2. We can also prove Lovász's result directly from Δ_1 using some properties of the Johnson graph defined below.

Definition 1. Johnson graph is a simple graph whose vertex set is the family of m -sets of an N -set. Two vertices U and V are adjacent if and only if $|U \cap V| = m - 1$.

Let J be the adjacency matrix of the Johnson graph with parameters $N = \binom{n}{2}$ and m . Let the square matrix B be defined as follows. The rows and columns of B are indexed by all the labelled m -edge graphs on a fixed set of n vertices, and ij -th entry is the number of ways of removing an edge from G_j and adding an edge to get G_i . Note that the diagonal entry is m , since we can add the same edge that is removed. Other entries of B are either 0 or 1. The matrix A is defined similarly for unlabelled graphs with m edges and n vertices. Thus matrix A is the matrix Δ_1 of the previous section. Matrix P is defined by indexing the rows by unlabelled graphs and columns by labelled graphs, and the ij -th entry is 1 if the labelled graph G_j is isomorphic to the unlabelled graph G_i . Other entries are 0. As in [ER], one can verify that $AP = PB$, and every eigenvalue of A is also an eigenvalue of B . But $B = mI + J$, and $-m$ is an eigenvalue of the Johnson graph if and only if $m \leq N/2$, (see [BCN]). So all eigenvalues of A are nonzero when $m > N/2$. This implies Lovász's result for any collection of graphs. Theorem 2.7 can then be proved from this, but we have included the first proof because it is constructive.

Perhaps even the k -edge version can be proved this way. In case of k -edge problem, B is simply $\sum_{i=0}^k \binom{m-i}{k-i} J_i$ where $J_0 = I$ and J_i is the matrix of the i -th relation of Johnson scheme. The eigenvalues of matrices in a Johnson algebra are known: the j -th eigenvalue ($j \leq \min(m, N - m)$) of B is

$$\sum_{i=0}^k \sum_{l=0}^i (-1)^l \binom{j}{l} \binom{m-j}{i-l} \binom{N-m-j}{i-l} \binom{m-i}{k-i}$$

We need to prove that when $2m - k + 1 > N$, above polynomial in j doesn't vanish in the range of integers 0 to $\min(m, N - m)$.

3. We have not solved the k -edge version of the problem of Krasikov and Roditty. But, since the procedure in the proof of Theorem 6 can be used also in the k -edge case, it would be enough to prove the k -edge deck version of Lovász's result.

To prove the k -edge deck version of Lovász's result we have indicated one approach using Johnson schemes. Another approach would be to relate Δ_i , $i \geq k$ to Δ_k explicitly, and then prove Lovász's result using those decks.

Acknowledgements

This work was done at Combinatorics and Optimization, University of Waterloo, Canada, and supported by the grant A7331 from NSERC. I take this opportunity to thank Adrian Bondy who arranged my visit. He and Hugh Hind were very helpful and encouraging throughout my stay at Waterloo. I would also like to thank the referees for their many valuable suggestions.

References

- [B] J.A. Bondy and R. L. Hemminger, Graph reconstruction - a survey, *J. Graph Theory* **1** no. 3 (1977) 227-268.
- [BCN] A. E. Brouwer, A. M. Cohen and A. Neumaier, *Distance-Regular Graphs*, Springer, Berlin, 1989.
- [ER] M. N. Ellingham and G. F. Royle, Vertex-switching reconstruction of subgraph numbers and triangle-free graphs, *J. Combinatorial Theory Ser. B* **54** no. 2 (1992) 167-177.
- [KR] I. Krasikov and Y. Roditty, Balance equations for reconstruction problems, *Arch. Math. (Basel)* **48** no. 5 (1987) 458-464.
- [L] L. Lovász, A note on the line reconstruction problem, *J. Combinatorial Theory Ser. B* **13** (1972) 309-310.
- [M] V. Müller, The edge reconstruction hypothesis is true for graphs with more than $n \cdot \log_2 n$ edges, *J. Combinatorial Theory Ser. B* **22** no. 3 (1977) 281-283.